

Signaling games with pattern recognition

Haoyang Wu

Department of Physics

Xi'an Jiaotong University, China.

hywch@mail.xjtu.edu.cn

Abstract

The classical model of signaling games assumes that the receiver exactly know the type space (private information) of the sender and be able to discriminate each type of the sender distinctly. However, the justification of this assumption is questionable. It is more reasonable to let the receiver recognize the pattern of the sender. In this paper, we investigate what happens if the assumption is relaxed. A framework of signaling games with pattern recognition and an example are given.

Keyword: Signaling games; pattern recognition.

1 Introduction

Since the pioneering work of Spence[1], signaling games have been deeply investigated by many researchers around the world [2-10]. Generally, there are two players in the model, a Sender (S) and a Receiver (R). The Sender has a certain type t (an element of a finite set T) which represents his private information. The Sender observes his own type while the Receiver does not know the type of the Sender. There is a strictly positive probability distribution $p(t)$ on T : $p(t)$ represents the prior probability that the Sender is of type t and is the common knowledge.

Based on his knowledge of his own type, the Sender chooses to

send a message from a set of possible messages $M = \{m_1, \dots, m_J\}$. The Receiver observes the message but not the type of the Sender. Then the Receiver chooses an action from a set of feasible actions $A = \{a_1, \dots, a_L\}$. The two players receive payoffs dependent on the Sender's type, the message chosen by the Sender and the action chosen by the Receiver.

Although the aforementioned model has been widely used by the economic community, people seldom notice clearly that they have made an assumption in the model, i.e., the Receiver is able to discriminate each type of the Sender distinctly. Actually, only by this assumption can the Receiver be able to have a probability distribution on the type set T . However, the justification of this assumption is questionable. Because the type space T is the private information of the Sender, the Sender has no incentive to tell his secret to the Receiver. The Receiver cannot take it for granted that he can discriminate each type of the Sender exactly.

The aim of this paper is to investigate what happens if the aforementioned assumption is canceled. We claim that what the Receiver can do is to recognize the “pattern” of the Sender (The phrase “pattern recognition” comes from computer science). A pattern consists of one type or some types, and one type just belongs to a pattern. The rest of the paper is organized as follows: Section 2 discusses the model of signaling games with pattern recognition. In Section 3, an example is given in detail.

2 The model

Without loss of generality, consider two players in a signaling game with pattern recognition, $i = 1, 2$. Player 1 is the Sender who has private information (i.e., the type). Player 2 is the Receiver. Different from classical signaling games (where the Receiver is assumed to know the type space of the sender and discriminate each type exactly), here we relax this assumption and

let the Receiver recognize the pattern of the sender (A message can be sent from several patterns. A pattern can send several messages). The pattern space and the belief distribution hold by the Receiver are common knowledge. The dynamic timing sequence of a signaling game with pattern recognition is as follows:

Step 1: Nature (N) selects the type t of player 1, $t \in T = \{t_1, \dots, t_K\}$, $K \geq 2$. Player 1 knows t , but player 2 doesn't know it. Furthermore, player 2 cannot discriminate each type in T exactly. The pattern space hold by player 2 is $T' = \{t'_1, \dots, t'_{\bar{K}}\}$, $1 \leq \bar{K} \leq K$. T' is a partition of T . The belief distribution hold by player 2 is defined on the pattern space T' : $p(t') > 0$, $t' \in T'$, $\sum_{t' \in T'} p(t') = 1$. A pattern t' consists of some types, and one type belongs to a pattern.

Step 2: After observing the type t , player 1 selects a message m from his message space $M = \{m_1, \dots, m_J\}$.

Step 3: After receiving the message m from player 1, player 2 makes an induction $p(t'|m)$ using Bayesian rules, and selects an action a from his feasible action space $A = \{a_1, \dots, a_L\}$.

Definition 1: The payoffs of player 1 and player 2 are both related to the type t of player 1, the message m and the action a of player 2, i.e., $u_1(t, m, a)$ and $u_2(t, m, a)$.

Note: 1) The strategy s_1 of player 1 is a message profile, which is related to the type t of player 1 and the pattern space T' hold by player 2, i.e., $s_1 = (m_{11}, \dots, m_{1K})$, where $m_{1k}(k, T') \in M = \{m_1, \dots, m_J\}$, $k = 1, \dots, K$.

2) The strategy s_2 of player 2 is an action profile, which is related to the message m and the pattern space T' hold by player 2, i.e., $s_2 = (a_{21}, \dots, a_{2J})$, where $a_{2j}(m, T') \in A = \{a_1, \dots, a_L\}$, $j = 1, \dots, J$.

As shown in Fig. 1, $K = 10$, $\bar{K} = 4$. $t'_1 = \{t_1, t_2, t_3\}$, $t'_2 = \{t_4, t_5, t_6, t_7\}$, $t'_3 = \{t_8\}$, $t'_4 = \{t_9, t_{10}\}$. To describe the perfect




Fig. 1. The type space and the pattern space.

Bayesian equilibrium of a signaling game with pattern recognition, we give five basic requirements as follows:

R1: The receiver has a belief about which patterns can have sent message m . These beliefs can be described as a probability distribution $p(t'|m)$, the probability that the sender is of pattern t' if the sender chooses message m . The sum over all patterns $t' \in T'$ of these probabilities has to be 1 conditional on any message m .

R2: The receiver can observe the probability distribution of messages, $p(m|t')$, when a pattern t' is given.

R3: The action that the receiver chooses must maximize the expected utility of the receiver given his beliefs about which patterns can have sent message m . This means that the sum $\sum_{t' \in T'} p(t'|m) \min_{t \in t'} u_2(t, m, a)$ is maximized. The action a that maximizes this sum is denoted as $a^*(m, T')$. For simplicity, we still use $a^*(m)$ to represent $a^*(m, T')$.

R4: For each type $t \in t'$ that the sender may have, the sender chooses to send the message

$$m^*(t') \in M^*(t') \equiv \{m \mid \max_{m \in M, t \in t'} u_1(t, m, a^*(m))\}$$

that maximizes the sender's utility $u_1(t, m, a^*(m))$ given the strategy chosen by the receiver $a^*(m)$.

R5: Let $|M^*(t')|$ be the cardinality of $M^*(t')$.

$$\tilde{p}(m|t') = \begin{cases} 1/|M^*(t')|, & m \in M^*(t') \\ 0, & \text{otherwise} \end{cases} \quad (1)$$

Let $T'(m)$ denote the set of patterns that have incentives to send m , i.e., $T'(m) = \{t' \in T' | m \in M^*(t')\}$. For each message m that the sender can send, if there exists a type $t \in t'$ such that $m^*(t')$ assigns strictly positive probability to m , the belief $p(t'|m)$ that the receiver has about the pattern t' of the sender if he observes the message m satisfies the Bayesian rule:

$$\tilde{p}(t'|m) = \frac{p(t')\tilde{p}(m|t')}{\sum_{t' \in T'(m)} p(t')\tilde{p}(m|t')} \quad (2)$$

Definition 2: A perfect Bayesian equilibrium of a signaling game with pattern recognition is composed of $m^*(t)$, $a^*(m)$, $p(t'|m)$ and $p(m|t')$, which satisfies R1–R5.

3 An example

Consider a car trade game. There are two players in the market, player 1 is the Seller, player 2 is the Buyer. The type space of player 1 is $T = \{t_1, t_2, t_3\}$, where t_1, t_2, t_3 represent low, medium and high quality of his car respectively. Player 1 knows his type but player 2 doesn't know it. The pattern space hold by player 2 is $T' = \{t'_1, t'_2\}$, $t'_1 = \{t_1, t_2\}$, $t'_2 = \{t_3\}$, i.e., in player 2's mind, there are two patterns: “Not high” and “High”. The belief distribution hold by player 2 is $p(t'_1) = \alpha$, $p(t'_2) = 1 - \alpha$.

The message space of player 1 is $M = \{m_1, m_2, m_3\}$, where m_1 , m_2 and m_3 represent three bids of the Seller, $m_3 > m_2 > m_1 > 0$. The action space of player 2 is $A = \{a_1, a_2\}$, where a_1 stands for “Not buy”, a_2 stands for “Buy”. The Seller announces his bids to the Buyer, and then the Buyer selects his action.

To describe the payoff functions of two players, $u_1(t, m, a)$ and $u_2(t, m, a)$, we assume that the cost for the Seller of type t_1 pretending to be of type t_2 is c_{12} . Similarly, we define c_{13} and c_{23} , and assume $c_{13} > c_{12} > 0$, $c_{13} > c_{23} > 0$. The value for the Buyer when he buy a car of type t_1, t_2, t_3 are V_1, V_2, V_3 respectively, $V_3 > V_2 > V_1 > 0$.

1) When the Buyer selects a_1 :

$$\begin{aligned} u_1(t_k, m_j, a_1) &= 0, k = 1 \sim 3, j = 1 \sim 3, j \leq k \\ u_1(t_1, m_2, a_1) &= -c_{12}, u_1(t_1, m_3, a_1) = -c_{13}, u_1(t_2, m_3, a_1) = -c_{23} \\ u_2(t_k, m_j, a_1) &= 0, k = 1 \sim 3, j = 1 \sim 3 \end{aligned}$$

2) When the Buyer selects a_2 :

$$\begin{aligned} u_1(t_1, m_1, a_2) &= m_1, & u_2(t_1, m_1, a_2) &= V_1 - m_1 \\ u_1(t_1, m_2, a_2) &= m_2 - c_{12}, & u_2(t_1, m_2, a_2) &= V_1 - m_2 \\ u_1(t_1, m_3, a_2) &= m_3 - c_{13}, & u_2(t_1, m_3, a_2) &= V_1 - m_3 \\ u_1(t_2, m_1, a_2) &= m_1, & u_2(t_2, m_1, a_2) &= V_2 - m_1 \\ u_1(t_2, m_2, a_2) &= m_2, & u_2(t_2, m_2, a_2) &= V_2 - m_2 \\ u_1(t_2, m_3, a_2) &= m_3 - c_{23}, & u_2(t_2, m_3, a_2) &= V_2 - m_3 \\ u_1(t_3, m_1, a_2) &= m_1, & u_2(t_3, m_1, a_2) &= V_3 - m_1 \\ u_1(t_3, m_2, a_2) &= m_2, & u_2(t_3, m_2, a_2) &= V_3 - m_2 \\ u_1(t_3, m_3, a_2) &= m_3, & u_2(t_3, m_3, a_2) &= V_3 - m_3 \end{aligned}$$

In order to obtain the perfect Bayesian equilibrium of the second-hand car trade game, we show the requirements R1–R5 in details:

R1: $\rho = p(t'_1|m_1)$, $1 - \rho = p(t'_2|m_1)$. $\sigma = p(t'_1|m_2)$, $1 - \sigma = p(t'_2|m_2)$. $\lambda = p(t'_1|m_3)$, $1 - \lambda = p(t'_2|m_3)$.

R2: Player 2 can observe the probability $p(m_j|t'_1)$ and $p(m_j|t'_2)$, $j = 1 \sim 3$.

R3: In order to computing the best strategy (action) $a^*(m)$ of player 2 when he receives a message $m \in M$, we need to solve

the optimization problem:

$$\max_{a \in A} \sum_{t' \in T'} p(t'|m) \min_{t \in t'} u_2(t, m, a) \quad (3)$$

For this example, the object function is represented as follows:

$$\begin{aligned} & p(t'_1|m) \min_{t \in t'_1} u_2(t, m, a) + p(t'_2|m) \min_{t \in t'_2} u_2(t, m, a) \\ &= p(t'_1|m) \min\{u_2(t_1, m, a), u_2(t_2, m, a)\} + p(t'_2|m) u_2(t_3, m, a) \end{aligned}$$

If the Buyer selects a_1 , the object function is 0. If the Buyer selects a_2 , the object function is:

$$p(t'_1|m) u_2(t_1, m, a) + p(t'_2|m) u_2(t_3, m, a) \quad (4)$$

1) If player 1 sends a message m_1 , formula (4) will be:

$$\rho(V_1 - m_1) + (1 - \rho)(V_3 - m_1) = (V_3 - m_1) - \rho(V_3 - V_1) \quad (5)$$

Comparing formula (5) with zero, there holds:

$$a^*(m_1) = \begin{cases} a_1, & \rho_0 \leq \rho \leq 1 \\ a_2, & 0 \leq \rho < \rho_0 \end{cases}, \text{ where } \rho_0 = \frac{V_3 - m_1}{V_3 - V_1}.$$

2) If player 1 sends a message m_2 , formula (4) will be:

$$\sigma(V_1 - m_2) + (1 - \sigma)(V_3 - m_2) = (V_3 - m_2) - \sigma(V_3 - V_1) \quad (6)$$

Comparing formula (6) with zero, there holds:

$$a^*(m_2) = \begin{cases} a_1, & \sigma_0 \leq \sigma \leq 1 \\ a_2, & 0 \leq \sigma < \sigma_0 \end{cases}, \text{ where } \sigma_0 = \frac{V_3 - m_2}{V_3 - V_1}.$$

3) If player 1 sends a message m_3 , formula (4) will be:

$$\lambda(V_1 - m_3) + (1 - \lambda)(V_3 - m_3) = (V_3 - m_3) - \lambda(V_3 - V_1) \quad (7)$$

Comparing formula (7) with zero, there holds:

$$a^*(m_3) = \begin{cases} a_1, & \lambda_0 \leq \lambda \leq 1 \\ a_2, & 0 \leq \lambda < \lambda_0 \end{cases}, \text{ where } \lambda_0 = \frac{V_3 - m_3}{V_3 - V_1}.$$

To sum the aforementioned optimal strategies of the Buyer, there will be the following eight kinds of outcomes:

1) For $(\rho, \sigma, \lambda) \in D_1 = \{\rho_0 \leq \rho \leq 1, \sigma_0 \leq \sigma \leq 1, \lambda_0 \leq \lambda \leq 1\}$,
 $a^*(m) = a_1, m \in M$.

2) For $(\rho, \sigma, \lambda) \in D_2 = \{\rho_0 \leq \rho \leq 1, \sigma_0 \leq \sigma \leq 1, 0 \leq \lambda < \lambda_0\}$,

$$a^*(m) = \begin{cases} a_1, & m = m_1, m_2 \\ a_2, & m = m_3 \end{cases}.$$

3) For $(\rho, \sigma, \lambda) \in D_3 = \{\rho_0 \leq \rho \leq 1, 0 \leq \sigma < \sigma_0, \lambda_0 \leq \lambda \leq 1\}$,

$$a^*(m) = \begin{cases} a_1, & m = m_1, m_3 \\ a_2, & m = m_2 \end{cases}.$$

4) For $(\rho, \sigma, \lambda) \in D_4 = \{\rho_0 \leq \rho \leq 1, 0 \leq \sigma < \sigma_0, 0 \leq \lambda < \lambda_0\}$,

$$a^*(m) = \begin{cases} a_1, & m = m_1 \\ a_2, & m = m_2, m_3 \end{cases}.$$

5) For $(\rho, \sigma, \lambda) \in D_5 = \{0 \leq \rho < \rho_0, \sigma_0 \leq \sigma \leq 1, \lambda_0 \leq \lambda \leq 1\}$,

$$a^*(m) = \begin{cases} a_1, & m = m_2, m_3 \\ a_2, & m = m_1 \end{cases}.$$

6) For $(\rho, \sigma, \lambda) \in D_6 = \{0 \leq \rho < \rho_0, \sigma_0 \leq \sigma \leq 1, 0 \leq \lambda < \lambda_0\}$,

$$a^*(m) = \begin{cases} a_1, & m = m_2 \\ a_2, & m = m_1, m_3 \end{cases}.$$

7) For $(\rho, \sigma, \lambda) \in D_7 = \{0 \leq \rho < \rho_0, 0 \leq \sigma < \sigma_0, \lambda_0 \leq \lambda \leq 1\}$,

$$a^*(m) = \begin{cases} a_1, & m = m_3 \\ a_2, & m = m_1, m_2 \end{cases}.$$

8) For $(\rho, \sigma, \lambda) \in D_8 = \{0 \leq \rho < \rho_0, 0 \leq \sigma < \sigma_0, 0 \leq \lambda < \lambda_0\}$,
 $a^*(m) = a_2, m \in M$.

R4: For each type t ($t \in t'$) that the Seller may have, we compute the optimal message $m^*(t') \in M^*(t')$ for the Seller in these eight zones $D_1, \dots, D_8$, i.e., we need to solve the maximization

problem:

$$\max_{m \in M, t \in T'} u_1(t, m, a^*(m)) \quad (8)$$

1) Consider the zone D_1 . There exists $a^*(m) = a_1, m \in M$.

For $t = t_1$,

$$\begin{aligned} u_1(t_1, m_1, a^*(m_1)) &= u_1(t_1, m_1, a_1) = 0 \\ u_1(t_1, m_2, a^*(m_2)) &= u_1(t_1, m_2, a_1) = -c_{12} \\ u_1(t_1, m_3, a^*(m_3)) &= u_1(t_1, m_3, a_1) = -c_{13} \end{aligned}$$

$$m^*(t_1) = m_1. \quad (9)$$

For $t = t_2$,

$$\begin{aligned} u_1(t_2, m_1, a^*(m_1)) &= u_1(t_2, m_1, a_1) = 0 \\ u_1(t_2, m_2, a^*(m_2)) &= u_1(t_2, m_2, a_1) = 0 \\ u_1(t_2, m_3, a^*(m_3)) &= u_1(t_2, m_3, a_1) = -c_{23} \end{aligned}$$

$$m^*(t_2) = m_1 \text{ or } m^*(t_2) = m_2. \quad (10)$$

For $t = t_3$,

$$\begin{aligned} u_1(t_3, m_1, a^*(m_1)) &= u_1(t_3, m_1, a_1) = 0 \\ u_1(t_3, m_2, a^*(m_2)) &= u_1(t_3, m_2, a_1) = 0 \\ u_1(t_3, m_3, a^*(m_3)) &= u_1(t_3, m_3, a_1) = 0 \end{aligned}$$

$$m^*(t_3) = m_1 \text{ or } m^*(t_3) = m_2 \text{ or } m^*(t_3) = m_3. \quad (11)$$

Comparing formula (9), (10) and (11), we obtain the optimal message for the Seller:

$$\text{For } (\rho, \sigma, \lambda) \in D_1, m^*(t) = m_1, t \in T; \text{ or } m^*(t) = \begin{cases} m_1, & t = t_1, t_2; \\ m_2, & t = t_3 \end{cases};$$

$$\text{or } m^*(t) = \begin{cases} m_1, & t = t_1, t_3; \\ m_2, & t = t_2 \end{cases}; \text{ or } m^*(t) = \begin{cases} m_1, & t = t_1 \\ m_2, & t = t_2, t_3 \end{cases}; \text{ or}$$

$$m^*(t) = \begin{cases} m_1, & t = t_1 \\ m_2, & t = t_2. \\ m_3, & t = t_3 \end{cases}$$

Therefore, for $(\rho, \sigma, \lambda) \in D_1, M^*(t') = \{m_1\}, t' \in T'; \text{ or } M^*(t') =$

$$\begin{aligned}
& \begin{cases} \{m_1\}, & t' = t'_1 \\ \{m_2\}, & t' = t'_2 \end{cases}, \\
& \text{or } M^*(t') = \begin{cases} \{m_1, m_2\}, & t' = t'_1 \\ \{m_1\}, & t' = t'_2 \end{cases}; \text{ or } M^*(t') = \begin{cases} \{m_1, m_2\}, & t' = t'_1 \\ \{m_2\}, & t' = t'_2 \end{cases}, \\
& \text{or } M^*(t') = \begin{cases} \{m_1, m_2\}, & t' = t'_1 \\ \{m_3\}, & t' = t'_2 \end{cases}.
\end{aligned}$$

$$2) \text{ Consider the zone } D_2. \text{ There exists } a^*(m) = \begin{cases} a_1, & m = m_1, m_2 \\ a_2, & m = m_3 \end{cases}.$$

For $t = t_1$,

$$\begin{aligned}
u_1(t_1, m_1, a^*(m_1)) &= u_1(t_1, m_1, a_1) = 0 \\
u_1(t_1, m_2, a^*(m_2)) &= u_1(t_1, m_2, a_1) = -c_{12} \\
u_1(t_1, m_3, a^*(m_3)) &= u_1(t_1, m_3, a_2) = m_3 - c_{13}
\end{aligned}$$

$$m^*(t_1) = \begin{cases} m_1, & c_{13} \geq m_3 \\ m_3, & 0 < c_{13} < m_3 \end{cases}. \quad (12)$$

For $t = t_2$,

$$\begin{aligned}
u_1(t_2, m_1, a^*(m_1)) &= u_1(t_2, m_1, a_1) = 0 \\
u_1(t_2, m_2, a^*(m_2)) &= u_1(t_2, m_2, a_1) = 0 \\
u_1(t_2, m_3, a^*(m_3)) &= u_1(t_2, m_3, a_2) = m_3 - c_{23}
\end{aligned}$$

$$m^*(t_2) = \begin{cases} m_1, & c_{23} \geq m_3 \\ m_3, & 0 < c_{23} < m_3 \end{cases}, \text{ or } m^*(t_2) = \begin{cases} m_2, & c_{23} \geq m_3 \\ m_3, & 0 < c_{23} < m_3 \end{cases}. \quad (13)$$

For $t = t_3$,

$$\begin{aligned}
u_1(t_3, m_1, a^*(m_1)) &= u_1(t_3, m_1, a_1) = 0 \\
u_1(t_3, m_2, a^*(m_2)) &= u_1(t_3, m_2, a_1) = 0 \\
u_1(t_3, m_3, a^*(m_3)) &= u_1(t_3, m_3, a_2) = m_3
\end{aligned}$$

$$m^*(t_3) = m_3. \quad (14)$$

Comparing formula (12), (13) and (14), we obtain the optimal message for the Seller:

For $(\rho, \sigma, \lambda) \in D_2$,

- If $0 < c_{13} < m_3$ and $0 < c_{23} < m_3$, then $m^*(t) = m_3, t \in T$.
- If $0 < c_{13} < m_3$ and $c_{23} \geq m_3$, then $m^*(t) = \begin{cases} m_1, & t = t_2 \\ m_3, & t = t_1, t_3 \end{cases}$,
or $m^*(t) = \begin{cases} m_2, & t = t_2 \\ m_3, & t = t_1, t_3 \end{cases}$.
- If $c_{13} \geq m_3$ and $0 < c_{23} < m_3$, then $m^*(t) = \begin{cases} m_1, & t = t_1 \\ m_3, & t = t_2, t_3 \end{cases}$.
- If $c_{13} \geq m_3$ and $c_{23} \geq m_3$, then $m^*(t) = \begin{cases} m_1, & t = t_1, t_2 \\ m_3, & t = t_3 \end{cases}$,
or $m^*(t) = \begin{cases} m_1, & t = t_1 \\ m_2, & t = t_2 \\ m_3, & t = t_3 \end{cases}$.

Therefore, for $(\rho, \sigma, \lambda) \in D_2$,

- If $0 < c_{13} < m_3$, then $M^*(t') = \{m_3\}, t' \in T'$.
- If $c_{13} \geq m_3$ and $0 < c_{23} < m_3$, then $M^*(t') = \{m_3\}, t' \in T'$.
- If $c_{13} \geq m_3$ and $c_{23} \geq m_3$, then $M^*(t') = \begin{cases} \{m_1\}, & t' = t'_1 \\ \{m_3\}, & t' = t'_2 \end{cases}$, or
 $M^*(t') = \begin{cases} \{m_1, m_2\}, & t' = t'_1 \\ \{m_3\}, & t' = t'_2 \end{cases}$.

3) Consider the zone D_3 . There exists $a^*(m) = \begin{cases} a_1, & m = m_1, m_3 \\ a_2, & m = m_2 \end{cases}$.

For $t = t_1$,

$$\begin{aligned} u_1(t_1, m_1, a^*(m_1)) &= u_1(t_1, m_1, a_1) = 0 \\ u_1(t_1, m_2, a^*(m_2)) &= u_1(t_1, m_2, a_2) = m_2 - c_{12} \\ u_1(t_1, m_3, a^*(m_3)) &= u_1(t_1, m_3, a_1) = -c_{13} \end{aligned}$$

$$m^*(t_1) = \begin{cases} m_1, & c_{12} \geq m_2 \\ m_2, & 0 < c_{12} < m_2 \end{cases}. \quad (15)$$

For $t = t_2$,

$$\begin{aligned} u_1(t_2, m_1, a^*(m_1)) &= u_1(t_2, m_1, a_1) = 0 \\ u_1(t_2, m_2, a^*(m_2)) &= u_1(t_2, m_2, a_2) = m_2 \\ u_1(t_2, m_3, a^*(m_3)) &= u_1(t_2, m_3, a_1) = -c_{23} \end{aligned}$$

$$m^*(t_2) = m_2. \quad (16)$$

For $t = t_3$,

$$\begin{aligned} u_1(t_3, m_1, a^*(m_1)) &= u_1(t_3, m_1, a_1) = 0 \\ u_1(t_3, m_2, a^*(m_2)) &= u_1(t_3, m_2, a_2) = m_2 \\ u_1(t_3, m_3, a^*(m_3)) &= u_1(t_3, m_3, a_1) = 0 \end{aligned}$$

$$m^*(t_3) = m_2. \quad (17)$$

Comparing formula (15), (16) and (17), we obtain the optimal message for the Seller:

For $(\rho, \sigma, \lambda) \in D_3$,

- If $0 < c_{12} < m_2$, then $m^*(t) = m_2$, $t \in T$.

- If $c_{12} \geq m_2$, then $m^*(t) = \begin{cases} m_1, & t = t_1 \\ m_2, & t = t_2, t_3 \end{cases}$.

Therefore, for $(\rho, \sigma, \lambda) \in D_3$, $M^*(t') = \{m_2\}$, $t' \in T'$.

4) Consider the zone D_4 . There exists $a^*(m) = \begin{cases} a_1, & m = m_1 \\ a_2, & m = m_2, m_3 \end{cases}$.

For $t = t_1$,

$$\begin{aligned} u_1(t_1, m_1, a^*(m_1)) &= u_1(t_1, m_1, a_1) = 0 \\ u_1(t_1, m_2, a^*(m_2)) &= u_1(t_1, m_2, a_2) = m_2 - c_{12} \\ u_1(t_1, m_3, a^*(m_3)) &= u_1(t_1, m_3, a_2) = m_3 - c_{13} \end{aligned}$$

$$m^*(t_1) = \begin{cases} m_1, & c_{12} \geq m_2, c_{13} \geq m_3 \\ m_2, & 0 < c_{12} < m_2, c_{13} - c_{12} \geq m_3 - m_2 \\ m_3, & 0 < c_{13} < m_3, c_{13} - c_{12} < m_3 - m_2 \end{cases} \quad (18)$$

For $t = t_2$,

$$\begin{aligned} u_1(t_2, m_1, a^*(m_1)) &= u_1(t_2, m_1, a_1) = 0 \\ u_1(t_2, m_2, a^*(m_2)) &= u_1(t_2, m_2, a_2) = m_2 \\ u_1(t_2, m_3, a^*(m_3)) &= u_1(t_2, m_3, a_2) = m_3 - c_{23} \end{aligned}$$

$$m^*(t_2) = \begin{cases} m_2, & c_{23} \geq m_3 - m_2 \\ m_3, & 0 < c_{23} < m_3 - m_2 \end{cases} \quad (19)$$

For $t = t_3$,

$$\begin{aligned} u_1(t_3, m_1, a^*(m_1)) &= u_1(t_3, m_1, a_1) = 0 \\ u_1(t_3, m_2, a^*(m_2)) &= u_1(t_3, m_2, a_2) = m_2 \\ u_1(t_3, m_3, a^*(m_3)) &= u_1(t_3, m_3, a_2) = m_3 \end{aligned}$$

$$m^*(t_3) = m_3. \quad (20)$$

Comparing formula (18), (19) and (20), we obtain the optimal message for the Seller:

For $(\rho, \sigma, \lambda) \in D_4$,

- If $c_{12} \geq m_2, c_{13} \geq m_3, c_{23} \geq m_3 - m_2$, then

$$m^*(t) = \begin{cases} m_1, & t = t_1 \\ m_2, & t = t_2 \\ m_3, & t = t_3 \end{cases} \quad M^*(t') = \begin{cases} \{m_2\}, & t' = t'_1 \\ \{m_3\}, & t' = t'_2 \end{cases}$$

- If $c_{12} \geq m_2, c_{13} \geq m_3, 0 < c_{23} < m_3 - m_2$, then

$$m^*(t) = \begin{cases} m_1, & t = t_1 \\ m_3, & t = t_2, t_3 \end{cases} \quad M^*(t') = \{m_3\}, t' \in T'.$$

- If $0 < c_{12} < m_2, c_{13} - c_{12} \geq m_3 - m_2, c_{23} \geq m_3 - m_2$, then

$$m^*(t) = \begin{cases} m_2, & t = t_1, t_2 \\ m_3, & t = t_3 \end{cases} \quad M^*(t') = \begin{cases} \{m_2\}, & t' = t'_1 \\ \{m_3\}, & t' = t'_2 \end{cases}$$

- If $0 < c_{12} < m_2, c_{13} - c_{12} \geq m_3 - m_2, 0 < c_{23} < m_3 - m_2$, then

$$m^*(t) = \begin{cases} m_2, & t = t_1 \\ m_3, & t = t_2, t_3 \end{cases}. \quad M^*(t') = \{m_3\}, \quad t' \in T'.$$

- If $0 < c_{13} < m_3$, $c_{13} - c_{12} < m_3 - m_2$, $c_{23} \geq m_3 - m_2$, then

$$m^*(t) = \begin{cases} m_2, & t = t_2 \\ m_3, & t = t_1, t_3 \end{cases}.$$

If $m_3 - m_2 \geq c_{13}$, then $M^*(t') = \{m_3\}$, $t' \in T'$; else $M^*(t') = \begin{cases} \{m_2\}, & t' = t'_1 \\ \{m_3\}, & t' = t'_2 \end{cases}$.

- If $0 < c_{13} < m_3$, $c_{13} - c_{12} < m_3 - m_2$, $0 < c_{23} < m_3 - m_2$, then

$$m^*(t) = m_3, \quad t \in T. \quad M^*(t') = \{m_3\}, \quad t' \in T'.$$

5) Consider the zone D_5 . There exists $a^*(m) = \begin{cases} a_1, & m = m_2, m_3 \\ a_2, & m = m_1 \end{cases}$.

For $t = t_1$,

$$\begin{aligned} u_1(t_1, m_1, a^*(m_1)) &= u_1(t_1, m_1, a_2) = m_1 \\ u_1(t_1, m_2, a^*(m_2)) &= u_1(t_1, m_2, a_1) = -c_{12} \\ u_1(t_1, m_3, a^*(m_3)) &= u_1(t_1, m_3, a_1) = -c_{13} \end{aligned}$$

$$m^*(t_1) = m_1. \quad (21)$$

For $t = t_2$,

$$\begin{aligned} u_1(t_2, m_1, a^*(m_1)) &= u_1(t_2, m_1, a_2) = m_1 \\ u_1(t_2, m_2, a^*(m_2)) &= u_1(t_2, m_2, a_1) = 0 \\ u_1(t_2, m_3, a^*(m_3)) &= u_1(t_2, m_3, a_1) = -c_{23} \end{aligned}$$

$$m^*(t_2) = m_1. \quad (22)$$

For $t = t_3$,

$$\begin{aligned} u_1(t_3, m_1, a^*(m_1)) &= u_1(t_3, m_1, a_2) = m_1 \\ u_1(t_3, m_2, a^*(m_2)) &= u_1(t_3, m_2, a_1) = 0 \\ u_1(t_3, m_3, a^*(m_3)) &= u_1(t_3, m_3, a_1) = 0 \end{aligned}$$

$$m^*(t_3) = m_1. \quad (23)$$

Comparing formula (21), (22) and (23), we obtain the optimal message for the Seller: For $(\rho, \sigma, \lambda) \in D_5$, $m^*(t) = m_1$, $t \in T$. $M^*(t') = \{m_1\}$, $t' \in T'$.

6) Consider the zone D_6 . There exists $a^*(m) = \begin{cases} a_1, & m = m_2 \\ a_2, & m = m_1, m_3 \end{cases}$.

For $t = t_1$,

$$\begin{aligned} u_1(t_1, m_1, a^*(m_1)) &= u_1(t_1, m_1, a_2) = m_1 \\ u_1(t_1, m_2, a^*(m_2)) &= u_1(t_1, m_2, a_1) = -c_{12} \\ u_1(t_1, m_3, a^*(m_3)) &= u_1(t_1, m_3, a_2) = m_3 - c_{13} \end{aligned}$$

$$m^*(t_1) = \begin{cases} m_1, & c_{13} \geq m_3 - m_1 \\ m_3, & 0 < c_{13} < m_3 - m_1 \end{cases}. \quad (24)$$

For $t = t_2$,

$$\begin{aligned} u_1(t_2, m_1, a^*(m_1)) &= u_1(t_2, m_1, a_2) = m_1 \\ u_1(t_2, m_2, a^*(m_2)) &= u_1(t_2, m_2, a_1) = 0 \\ u_1(t_2, m_3, a^*(m_3)) &= u_1(t_2, m_3, a_2) = m_3 - c_{23} \end{aligned}$$

$$m^*(t_2) = \begin{cases} m_1, & c_{23} \geq m_3 - m_1 \\ m_3, & 0 < c_{23} < m_3 - m_1 \end{cases}. \quad (25)$$

For $t = t_3$,

$$\begin{aligned} u_1(t_3, m_1, a^*(m_1)) &= u_1(t_3, m_1, a_2) = m_1 \\ u_1(t_3, m_2, a^*(m_2)) &= u_1(t_3, m_2, a_1) = 0 \\ u_1(t_3, m_3, a^*(m_3)) &= u_1(t_3, m_3, a_2) = m_3 \end{aligned}$$

$$m^*(t_3) = m_3. \quad (26)$$

Comparing formula (24), (25) and (26), we obtain the optimal message for the Seller:

For $(\rho, \sigma, \lambda) \in D_6$,

- If $c_{23} \geq m_3 - m_1$, then $m^*(t) = \begin{cases} m_1, & t = t_1, t_2 \\ m_3, & t = t_3 \end{cases}$.

- If $0 < c_{23} < m_3 - m_1 \leq c_{13}$, then $m^*(t) = \begin{cases} m_1, & t = t_1 \\ m_3, & t = t_2, t_3 \end{cases}$.
- If $0 < c_{13} < m_3 - m_1$, then $m^*(t) = m_3, t \in T$.

Therefore, for $(\rho, \sigma, \lambda) \in D_6$,

- If $c_{23} \geq m_3 - m_1$, then $M^*(t') = \begin{cases} \{m_1\}, & t' = t'_1 \\ \{m_3\}, & t' = t'_2 \end{cases}$.
- If $0 < c_{23} < m_3 - m_1$, then $M^*(t') = \{m_3\}, t' \in T'$.

7) Consider the zone D_7 . There exists $a^*(m) = \begin{cases} a_1, & m = m_3 \\ a_2, & m = m_1, m_2 \end{cases}$.

For $t = t_1$,

$$\begin{aligned} u_1(t_1, m_1, a^*(m_1)) &= u_1(t_1, m_1, a_2) = m_1 \\ u_1(t_1, m_2, a^*(m_2)) &= u_1(t_1, m_2, a_2) = m_2 - c_{12} \\ u_1(t_1, m_3, a^*(m_3)) &= u_1(t_1, m_3, a_1) = -c_{13} \end{aligned}$$

$$m^*(t_1) = \begin{cases} m_1, & c_{12} \geq m_2 - m_1 \\ m_2, & 0 < c_{12} < m_2 - m_1 \end{cases}. \quad (27)$$

For $t = t_2$,

$$\begin{aligned} u_1(t_2, m_1, a^*(m_1)) &= u_1(t_2, m_1, a_2) = m_1 \\ u_1(t_2, m_2, a^*(m_2)) &= u_1(t_2, m_2, a_2) = m_2 \\ u_1(t_2, m_3, a^*(m_3)) &= u_1(t_2, m_3, a_1) = -c_{23} \end{aligned}$$

$$m^*(t_2) = m_2. \quad (28)$$

For $t = t_3$,

$$\begin{aligned} u_1(t_3, m_1, a^*(m_1)) &= u_1(t_3, m_1, a_2) = m_1 \\ u_1(t_3, m_2, a^*(m_2)) &= u_1(t_3, m_2, a_2) = m_2 \\ u_1(t_3, m_3, a^*(m_3)) &= u_1(t_3, m_3, a_1) = 0 \end{aligned}$$

$$m^*(t_3) = m_2. \quad (29)$$

Comparing formula (27), (28) and (29), we obtain the optimal message for the Seller:

For $(\rho, \sigma, \lambda) \in D_7$,

- If $c_{12} \geq m_2 - m_1$, then $m^*(t) = \begin{cases} m_1, & t = t_1 \\ m_2, & t = t_2, t_3 \end{cases}$.
- If $0 < c_{12} < m_2 - m_1$, then $m^*(t) = m_2, t \in T$.

Therefore, for $(\rho, \sigma, \lambda) \in D_7$, $M^*(t') = \{m_2\}, t' \in T'$.

8) Consider the zone D_8 . There exists $a^*(m) = a_2, m \in M$.

For $t = t_1$,

$$\begin{aligned} u_1(t_1, m_1, a^*(m_1)) &= u_1(t_1, m_1, a_2) = m_1 \\ u_1(t_1, m_2, a^*(m_2)) &= u_1(t_1, m_2, a_2) = m_2 - c_{12} \\ u_1(t_1, m_3, a^*(m_3)) &= u_1(t_1, m_3, a_2) = m_3 - c_{13} \end{aligned}$$

$$m^*(t_1) = \begin{cases} m_1, & c_{12} \geq m_2 - m_1, c_{13} \geq m_3 - m_1 \\ m_2, & 0 < c_{12} < m_2 - m_1, c_{13} - c_{12} \geq m_3 - m_2 \\ m_3, & 0 < c_{13} < m_3 - m_1, c_{13} - c_{12} < m_3 - m_2 \end{cases}. \quad (30)$$

For $t = t_2$,

$$\begin{aligned} u_1(t_2, m_1, a^*(m_1)) &= u_1(t_2, m_1, a_2) = m_1 \\ u_1(t_2, m_2, a^*(m_2)) &= u_1(t_2, m_2, a_2) = m_2 \\ u_1(t_2, m_3, a^*(m_3)) &= u_1(t_2, m_3, a_2) = m_3 - c_{23} \end{aligned}$$

$$m^*(t_2) = \begin{cases} m_2, & c_{23} \geq m_3 - m_2 \\ m_3, & 0 < c_{23} < m_3 - m_2 \end{cases}. \quad (31)$$

For $t = t_3$,

$$\begin{aligned} u_1(t_3, m_1, a^*(m_1)) &= u_1(t_3, m_1, a_2) = m_1 \\ u_1(t_3, m_2, a^*(m_2)) &= u_1(t_3, m_2, a_2) = m_2 \\ u_1(t_3, m_3, a^*(m_3)) &= u_1(t_3, m_3, a_2) = m_3 \end{aligned}$$

$$m^*(t_3) = m_3. \quad (32)$$

Comparing formula (30), (31) and (32), we obtain the optimal message for the Seller:

For $(\rho, \sigma, \lambda) \in D_8$,

- If $c_{12} \geq m_2 - m_1$, $c_{13} \geq m_3 - m_1$, $c_{23} \geq m_3 - m_2$, then

$$m^*(t) = \begin{cases} m_1, & t = t_1 \\ m_2, & t = t_2 \\ m_3, & t = t_3 \end{cases}. M^*(t') = \begin{cases} \{m_2\}, & t' = t'_1 \\ \{m_3\}, & t' = t'_2 \end{cases}.$$

- If $c_{12} \geq m_2 - m_1$, $c_{13} \geq m_3 - m_1$, $0 < c_{23} < m_3 - m_2$, then

$$m^*(t) = \begin{cases} m_1, & t = t_1 \\ m_3, & t = t_2, t_3 \end{cases}. M^*(t') = \{m_3\}, t' \in T'.$$

- If $0 < c_{12} < m_2 - m_1$, $c_{13} - c_{12} \geq m_3 - m_2$, $c_{23} \geq m_3 - m_2$, then

$$m^*(t) = \begin{cases} m_2, & t = t_1, t_2 \\ m_3, & t = t_3 \end{cases}. M^*(t') = \begin{cases} \{m_2\}, & t' = t'_1 \\ \{m_3\}, & t' = t'_2 \end{cases}.$$

- If $0 < c_{12} < m_2 - m_1$, $c_{13} - c_{12} \geq m_3 - m_2$, $0 < c_{23} < m_3 - m_2$, then

$$m^*(t) = \begin{cases} m_2, & t = t_1 \\ m_3, & t = t_2, t_3 \end{cases}. M^*(t') = \{m_3\}, t' \in T'.$$

- If $0 < c_{13} < m_3 - m_1$, $c_{13} - c_{12} < m_3 - m_2$, $c_{23} \geq m_3 - m_2$, then

$$m^*(t) = \begin{cases} m_2, & t = t_2 \\ m_3, & t = t_1, t_3 \end{cases}. M^*(t') = \begin{cases} \{m_2\}, & t' = t'_1 \\ \{m_3\}, & t' = t'_2 \end{cases}.$$

- If $0 < c_{13} < m_3 - m_1$, $c_{13} - c_{12} < m_3 - m_2$, $0 < c_{23} < m_3 - m_2$, then

$$m^*(t) = m_3, t \in T. M^*(t') = \{m_3\}, t' \in T'.$$

R5:

For $M^*(t') = \{m_1\}$, $t' \in T'$:

$$\begin{aligned} T'(m_1) &= \{t'_1, t'_2\}, T'(m_2) = T'(m_3) = \phi \\ \tilde{p}(m_1|t') &= 1, \tilde{p}(m_2|t') = \tilde{p}(m_3|t') = 0, t' \in T' \\ \tilde{p}(t'_1|m_1) &= \frac{p(t'_1)\tilde{p}(m_1|t'_1)}{p(t'_1)\tilde{p}(m_1|t'_1) + p(t'_2)\tilde{p}(m_1|t'_2)} = \alpha \end{aligned}$$

For $M^*(t') = \{m_2\}$, $t' \in T'$:

$$\begin{aligned} T'(m_1) &= T'(m_3) = \phi, T'(m_2) = \{t'_1, t'_2\} \\ \tilde{p}(m_1|t') &= \tilde{p}(m_3|t') = 0, \tilde{p}(m_2|t') = 1, t' \in T' \\ \tilde{p}(t'_1|m_2) &= \frac{p(t'_1)\tilde{p}(m_2|t'_1)}{p(t'_1)\tilde{p}(m_2|t'_1) + p(t'_2)\tilde{p}(m_2|t'_2)} = \alpha \end{aligned}$$

For $M^*(t') = \{m_3\}$, $t' \in T'$:

$$\begin{aligned} T'(m_1) &= T'(m_2) = \phi, T'(m_3) = \{t'_1, t'_2\} \\ \tilde{p}(m_1|t') &= \tilde{p}(m_2|t') = 0, \tilde{p}(m_3|t') = 1, t' \in T' \\ \tilde{p}(t'_1|m_3) &= \frac{p(t'_1)\tilde{p}(m_3|t'_1)}{p(t'_1)\tilde{p}(m_3|t'_1) + p(t'_2)\tilde{p}(m_3|t'_2)} = \alpha \end{aligned}$$

For $M^*(t') = \begin{cases} \{m_1\}, & t' = t'_1. \\ \{m_2\}, & t' = t'_2. \end{cases}$

$$\begin{aligned} T'(m_1) &= \{t'_1\}, T'(m_2) = \{t'_2\}, T'(m_3) = \phi \\ \tilde{p}(m_1|t'_1) &= \tilde{p}(m_2|t'_2) = 1, \tilde{p}(t'_1|m_1) = \tilde{p}(t'_2|m_2) = 1. \end{aligned}$$

For $M^*(t') = \begin{cases} \{m_1\}, & t' = t'_1. \\ \{m_3\}, & t' = t'_2. \end{cases}$

$$\begin{aligned} T'(m_1) &= \{t'_1\}, T'(m_2) = \phi, T'(m_3) = \{t'_2\} \\ \tilde{p}(m_1|t'_1) &= \tilde{p}(m_3|t'_2) = 1, \tilde{p}(t'_1|m_1) = \tilde{p}(t'_2|m_3) = 1. \end{aligned}$$

For $M^*(t') = \begin{cases} \{m_2\}, & t' = t'_1. \\ \{m_3\}, & t' = t'_2. \end{cases}$

$$\begin{aligned} T'(m_1) &= \phi, T'(m_2) = \{t'_1\}, T'(m_3) = \{t'_2\} \\ \tilde{p}(m_2|t'_1) &= \tilde{p}(m_3|t'_2) = 1, \tilde{p}(t'_1|m_2) = \tilde{p}(t'_2|m_3) = 1. \end{aligned}$$

$$\text{For } M^*(t') = \begin{cases} \{m_1, m_2\}, & t' = t'_1 \\ \{m_1\}, & t' = t'_2 \end{cases}.$$

$$\begin{aligned} T'(m_1) &= \{t'_1, t'_2\}, T'(m_2) = \{t'_1\}, T'(m_3) = \phi \\ \tilde{p}(m_1|t'_1) &= \tilde{p}(m_2|t'_1) = 0.5, \tilde{p}(m_1|t'_2) = 1, \\ \tilde{p}(t'_1|m_1) &= \frac{p(t'_1)\tilde{p}(m_1|t'_1)}{p(t'_1)\tilde{p}(m_1|t'_1) + p(t'_2)\tilde{p}(m_1|t'_2)} = \frac{\alpha}{2 - \alpha} \\ \tilde{p}(t'_2|m_1) &= \frac{2 - 2\alpha}{2 - \alpha}, \tilde{p}(t'_1|m_2) = 1. \end{aligned}$$

$$\text{For } M^*(t') = \begin{cases} \{m_1, m_2\}, & t' = t'_1 \\ \{m_2\}, & t' = t'_2 \end{cases}.$$

$$\begin{aligned} T'(m_1) &= \{t'_1\}, T'(m_2) = \{t'_1, t'_2\}, T'(m_3) = \phi \\ \tilde{p}(m_1|t'_1) &= \tilde{p}(m_2|t'_1) = 0.5, \tilde{p}(m_2|t'_2) = 1, \\ \tilde{p}(t'_1|m_2) &= \frac{p(t'_1)\tilde{p}(m_2|t'_1)}{p(t'_1)\tilde{p}(m_2|t'_1) + p(t'_2)\tilde{p}(m_2|t'_2)} = \frac{\alpha}{2 - \alpha}. \\ \tilde{p}(t'_2|m_2) &= \frac{2 - 2\alpha}{2 - \alpha}, \tilde{p}(t'_1|m_1) = 1. \end{aligned}$$

$$\text{For } M^*(t') = \begin{cases} \{m_1, m_2\}, & t' = t'_1 \\ \{m_3\}, & t' = t'_2 \end{cases}.$$

$$\begin{aligned} T'(m_1) &= T'(m_2) = \{t'_1\}, T'(m_3) = \{t'_2\} \\ \tilde{p}(m_1|t'_1) &= \tilde{p}(m_2|t'_1) = 0.5, \tilde{p}(m_3|t'_2) = 1, \\ \tilde{p}(t'_1|m_1) &= \tilde{p}(t'_1|m_2) = \tilde{p}(t'_2|m_3) = 1. \end{aligned}$$

References

- [1] Spence, M., 1973. Job market signaling. *Quart. J. Econ.* 87, 355-374.
- [2] Cho, I.K., Kreps, D., 1987. Signaling games and stable equilibria. *Quart. J. Econ.* 56, 601-612.
- [3] Banks, J.S., Sobel, J., 1987. Equilibrium selection in signaling games. *Econometrica* 55, 647-661.
- [4] Cho, I.K., Sobel, J., 1990. Strategic stability and uniqueness in signaling games. *J. Econ. Theory* 50, 381-413.

- [5] Mailath, G.J., Fujiwara, M.O., Postlewaite, A., 1993. Belief-based refinements in signaling games. *J. Econ. Theory* 60, 241-276.
- [6] Manelli, A.M., 1996. Cheap talk and sequential equilibria in signaling games. *Econometrica* 64, 917-942.
- [7] Stamland, T., 1999. Partially informative signaling. *J. Econ. Theory* 89, 148-161.
- [8] Riley, J.G., 2001. Silver signals: Twenty-five years of screening and signaling. *J. Econ. Lit.* 39, 432-478.
- [9] Kremer, I., Skrzypacz, A., 2007. Dynamic signaling and market breakdown. *J. Econ. Theory* 133, 58-82.
- [10] Kaya, A., 2009. Repeated signaling games. *Game Econ. Behav.* 66, 841-854.